



**THE EFFECT OF ANTENATAL CARE ON THE HEALTH OF PREGNANT WOMEN  
(STUDY IN THE PRIMARY HEALTH CENTER IN A NEW BAQUBAH CITY)**

**Dr. Majida Hameed\***

College of medicine university of Diyala.

**\*Corresponding Author: Dr. Majida Hameed**

College of medicine university of Diyala.

Article Received on 21/03/2018

Article Revised on 11/04/2018

Article Accepted on 02/05/2018

**ABSTRACT**

The study group consisted of a sample of files of pregnant women attending for antenatal care during 1993 and 1994 years. This study started at (1) of July and ended at first of September 1995 year. No. of sample were (200) files. Detailed information sheet (backgrounds factors, sociocultural factors and medical factors) was designed together the required information. We found in this study that women whose age was between (15 – 29 years) were the most frequent attenders of the antenatal clinic. As regards parity, Nulliparus was found to be the most frequent attenders at the antenatal clinic were most frequently read / write and followed by primary school graduates (45%) and (21%) respectively. Amount of prenatal care received can be determined by 2 parameters:- (I) Trimester of gestational when A.N.C was commenced. In this study (71.5%) of women attended antenatal clinic during the first trimester, and (23.5%) of women attended A.N.C at third trimester. (II) Second parameter is the No. of antenatal visits. Most of women attending the A.N. clinic had 7 – 10 antenatal visit (51%).

**KEYWORDS:** antenatal care, Nulliparus, prenatal, A.N.C.

**INTRODUCTION**

Women are important to P.H.C in three ways:-

1. As direct beneficiaries.
2. As mother and family members.
3. And as front line health workers outside home.

All these roles especially the first must be emphasized in order to reduce morbidity and mortality in developing countries. Women carry much of the responsibility for world producing in the family, on the farm, in both traditional and modern – sector industries and in commerce.

An estimated 18 to 30% of the world's families are solely supported by women (Uncief – 1986).

Maternal health is a prime determinant of infant health which are so interlinked that the promotion of one entails consideration of promotion of the other (W.H.O – 1991).

**AIMS OF THE STUDY**

The aims of this study is; to assess the effect of antenatal care on the health of pregnant women attending the primary health center in a new Baqubah city and to study activities of this center.

And to relate all these findings to certain demographic variables in the women studied.

**LITERATURE REVIEW**

The child and mother welfare movement started in Europe at the end of the nineteenth century at the beginning of the industrial revolution (Jawadi, 1973).

In France infant consultation clinics were established in (1892).

In Belgium another different approach to the problem was made, efforts directed towards educating the mothers. in general aspect relating to child care (Jawadi, 1973).

In England, women volunteers tried to save life of pregnant women by house visits (Jawadi, 1973).

The first antenatal clinic in British was opened in 1910.

**Factors affecting attendants of A.N. care**

- Background factors like age, race, marital status and parity.
- Social class of the family and their income.
- Occupation of mothers and husband.

**Back ground factor**

- A) Age:- Non registered mothers in A.N.C. tended to be older than registered mothers (Klein, 1971)
- B) Parity:- Nullipara attend early as a result of general anxiety about the unknown and to ensure hospital bed.

**Sociocultural factors****A- Education**

Women of higher education level have more examination and intervention Hubert et al., (1987)

**B- Employment status**

The employment status acts as an index of educational status i.e are more aware of the desirability of early attendance (Kaliszas and Kidd, 1981).

**C- Income and medical insurance**

Moore et al., (1986) found that there is a markedly higher financial cost of delivering a patient without prenatal care when compared with that of the patients given care.

**D- Attitude towards health professionals**

Gardiner (1969) found that an important factor contributing to inadequacy of antepartum visit in some of women was not the lack of quality of medical care, but the lack of understanding and appreciation on the part of health team of coexistence of major personal and social problem which the women considers of great current importance than any problem relating to the pregnancy.

**Structural factors**

- A- Accessibility to health service.
- B- Relation of distance to missed visits.

Russel, (1977) → There is some evidence supporting this hypothesis.

**Amount and timing of antenatal care**

These are determined by the month of the gestation that care begins, No. of visits and length of gestation.

The frequency of visits increased with increasing educational level, residence in an urban environment, presence of adverse outcome of previous pregnancy, and early occurrence of complication (Hubret et al., 1987).

Late attendance for antenatal care is more frequent among working class women and at the extremes of maternal age and parity. It is also associated with distance from the clinic.

**Responsibilities of M.C.H service in health center and Intra structures**

Monitoring the body weight of all pregnant women to assess gain in pregnant women and adopting appropriate measures whenever necessary:

1. For mother.
2. Re organization of health care service.

- a) Strengthening MCH in P.H.C.
- b) Active school health services, and school health education programmes.
- c) Provision of food supplements.
- d) Tetanus Toxoid inoculation as routine precaution for all pregnant women.
- e) Access to essential obstetrical care at all levels.

**Basic maternal health services**

- 1- Family planning education.
- 2- Nutrition education.
- 3- Home gardens.
- 4- Tetanus immunization.
- 5- Iodine injection (for goiter).

**Screening and the at – risk concept**

Screening pregnant women for the problems is an important step towards treating and preventing those problems.

Screening required identifying pregnant women and list of risk factors, trained personnel to screen, record system.

High risk women would need more frequent contacts.

**Typical risk factors: - (Uncief, 1984)**

1. Weight of 38 kg or less before pregnancy.
2. Weight of 42 or less at 34<sup>th</sup> week of pregnancy.
3. Height of less than 145 cm.
4. Sever pallor.
5. Child from previous delivery who weighed less than 2kg at birth.
6. High blood pressure.
7. Age before 18 or about 30 at first pregnancy.
8. Age 35 years or older at the time of pregnancy.
9. History of abortion / or stillbirth.
10. Fifth or later child.
11. History of bleeding during pregnancy.
12. Possible twins.
13. Previous C/S, and malpresentation.
14. Hepatitis.

**Identification of the patient at risk of preterm labour**

All preterm labour happen in nulliporous patient (Arias,1984).

**Factors associated with preterm labour and delivery are the following****1- Maternal characteristic**

- a) Weight of mother before pregnancy if less than 50kg have 3 times as many low birth weight infant, as a mother who weighs more than 57 kg before pregnancy (Brown et al, 1981).
- b) Smoking during pregnancy. (twice frequent) Reborn and Meyerm, 1978).
- c) Maternal age.

15.8% of small babies are born to mothers whose age is under 15 years (Arias, 1984).

**2- Obstetric factors**

- a- Maternal weight gain.
  - b- Previous spontaneous or induced abortion especially mid trimester.
  - c- Multiple pregnancy.
  - d- History of stillbirth or neonatal death in prior pregnancy.
- 3- Maternal disease.** e.g. eclampsia, preclampsia, chronic renal disease, chronic hypertension associated with low birth weight.
- 4- Fetal problems** and placental factors. also cause low birth weight.

**Nutrition during pregnancy**

Health education about nutrition is very important thing so health education of the pregnant women who attend antenatal clinic about nutrition is a must.

Over weight is associated frequently with perinatal problems as prolonged labour, post partum haemorrhage and infection and poor perinatal outcome are all more frequent in obese pregnant patient.

**MATERIALS AND METHODS**

This study was carried out at the primary health center of a new Baqubah city, Training center on the files of 200 pregnant women.

This study started from first of July to first of September 1995.

The study group consisted of women who received antenatal care during the years 1993 – 1994. Their records were selected from the record department in the primary health center in a new Baqubah city randomly in a systematic way. (Sample, frame).

A detailed information sheet including information on the followings was used during the study.

1. Age (years): The age of pregnant women was subdivided into 6 groups with 5 years intervals.
2. Residence.
3. Educational status of mothers and their husbands.
4. Past obstetric history.
5. Parity of pregnant women who attended antenatal clinic.

Information about attending antenatal clinic was taken considering the following parameters:

1. No. of antenatal visits.
2. Uterus size at first antenatal visit was subdivided as first trimester, second trimester and third trimester.
3. Weight gain during antenatal care.
4. Information about vaccination of pregnant women.
5. Information about hematinic supplementations.
6. Complete information about investigations such as blood group, G.U.E, random blood sugar for D.M. and V.D.R.L for syphilis.
7. Blood pressure during antenatal care.

**RESULTS****Master table (1)**

Shows that women whose age was between 15 – 29 years were the most frequent attenders.

**Table (2)**

Shows the relationship between age of pregnant women and trimester of first antenatal visits.

We can see that the younger women (20 – 24 y.) visited the antenatal care generally earlier than older women.

**Table (3)**

Shows the relationship between parity and No. of antenatal visits of pregnant women attending antenatal clinic.

We see that the No. of antenatal visit appeared to be higher in (7 – 10) visits in the primiparous women.

**Table (4)**

Shows the relationship between age of pregnant women and No. of antenatal visit.

- The younger age group < 20 years a higher percentage of mothers was detected in less than 3 visits.
- Higher percentage of 3 - 6 visits was detected in (21 – 24) years pregnant women. and also higher percentage of 7 – 10 visits was detected in pregnant women with age of (21 – 24) years. (23.52%).

**Table 1: Master table illustrating the sample description of all variables included in our study (in percentage).**

Age (yrs)	15-19	15-19	15-19	15-19	15-19	15-19	Total
	21	21	21	21	21	21	200
Parity	0	1-3	4-6	7+	Total		
	55	24	13	4	200		
Education of mother	illuturate Read write Primary Intermediat and 2° College						Total
	10	45		21	14	10	200
Education of father	4 40 14 17 25						200
Weight gain /m.	No gain 6kg 7-10kg 11-13kg 14kg						Total
	10	20.5	34.5	14.5	15.5		200
No. of A. N. C.	2 visits 3 0 6 7-10 11+						Total
	30.5%	52%	15.5%	1%			200
T.T vaccination of preg.women	1st+2nd dose of T.T 3rd dose of vaccin 1st does only No vaccin						Total
	55%		6%		15%	24%	200
Hematin supplementation	Tonie No supplementation						
	100% 0						
Postnatal checkup	Postnatal checking No checking						
	57%			43%			

**Table 2: Relationship between age of pregnant women and trimster of first antenatal visit.**

Age	No. % first trimester	No. % Second trimester	No. % Tird trimester	Total
15 - 19 years	32 22.37	8 17.02	4 40	44
20 - 24 years	45 31.46	19 40.42	2 20	66
25 - 29 years	34 23.77	9 1.91	2 20	45
30 - 34 years	17 11.88	5 1.06	1 10	23
35 - 39 years	12 8.39	6 1.27	1 10	19
40 +	3 2.09	0	0	23
	143 100	47 100	10 100	200

**Table 3: Relationship between parity and number of antenatal visits of pregnant women attending antenatal clinic.**

parity	< 3 No. %	3 - 6 No. %	7 - 10 No. %	11 + No. %	Total
0	28 46.6	63 61.76	18 54.5	0	109
1 - 3	19 31.6	25 24.5	4 12.1	1 2.0	49
4 - 6	10 31.6	10 9.8	6 18.18	0	26
7 +	3 5.2	4 3.92	5 15.15	4 25	16
Total	60 100	102 100	33 100	5	200

**Table 4: Relationship between age of pregnant women and No. of antenatal visit.**

Age	Number of antenatal Visit				Total
	< 3 No. %	3 - 6 No. %	7 - 10 No. %	11 + No. %	
15 - 19	18 31.03	26 24.52	6 17.64	1 1.96	51
21 - 24	15 25.86	30 28.30	8 23.52	1 1.85	54
25 - 29	14 24.13	22 20.75	7 20.58	0	43
30 - 34	5 8.62	10 9.43	5 14.70	0	20
35 - 39	4 6.89	10 9.43	3 8.82	0	17
40 +	2 3.44	8 7.54	5 14.70	0	15
	58 100	106 100	34 100	2	200

## DISCUSSION

This study was undertaken to see the effect of antenatal care on the health of pregnant women, and also we classify here the factors that may be related with receiving antenatal care into four main classes, these are:

- Background factors.
- Socio cultural factors.
- Medical factors.
- Amount of A.N. care received.

### 1) Background Factors

Age: age of pregnant women attending the A.N. clinic.

Table (4) shows that younger women tend to receive adequate ANC more frequently than older women.

This result is compatible with the studies conducted by Klein, 1971, Poland et al., 1987 who suggested that younger women attended A.N. clinic more frequently than older women.

Parity: In this study we see that primigravida had higher percentage of antenatal visit than multipara. So parity appeared to be related to the No. of antenatal visits, were more prone to ask for antenatal care than those who had multiple previous pregnancies.

The result of this study are supported by Kalizer and Kidd, 1981 who found that missed visits to antenatal clinic is at minimum in the multipara as well as the finding of Poland et al., 1987 who mentioned that there was significant difference among different parity group with better care receive by those with lower parity.

Higher No. of visits mean that the women is better motivate and take more iron supplementations more than those with low visits.

### 2) Sociocultural Factors

Education: In this study we see mothers of lower educational levels were most likely to have visited the primary health center in a new Baqubah city, than mothers with high educational status because the last more likely to have received care from private physician. In this study 45% of pregnant women who attend the center.

### 3) Medical Factors

Previous obstetrical history:

1. Women with a bad obstetrical history had better attendance to antenatal clinic. These results agreement with those of Bruce et al., 1979 study and Saleem (1981).
2. Weight gain during antenatal care:

In this study high frequency of suboptimal weight gain (34.5%), 7 – 10kg this frequency may be due to low socioeconomic status and crowded area and to short spacing between pregnancies, This leads to maternal depletion where no time is available to recover adequately from last pregnancy (Unicef, 1984).

### 4) Amount of prenatal care received

1. Month of gestational age that care begins can determine the amount of prenatal care received, (Kesses et al., 1973).

Some women believe that when the fetus moved they felt alright and didn't need to seek antenatal care. (Poland et al. 1987).

2. Number of antenatal visits:

This No. varies from 1 -14 visits, (Hubret, 1987).

The No. of A.N. visits increases with higher education, residence in urban environment, presence of adverse outcome of previous pregnancy and early occurrence of complication. (Hubert, et al., 1987).

### 5) Others

1. Vaccination: 95% coverage with tetanus toxoid is effective in preventing maternal and neonatal tetanus if given in 2 doses with boosters every 5 or 10 years. (Unicef, 1984).

The one million infant deaths each year caused by tetanus neonatorum could be prevented through proper immunization of the mother (before or during pregnancy) or care with hygiene during or after birth, (Unicef, 1984 / 1986).

This sample appeared to have a coverage with TT in good percentage.

1. Hematinic supplementations

Our study shows that 100% of women studied took hematinic supplementation. In order to reduce, the incidence of anemia.

**Postnatal check up:** 57% postnatal checking in this study.

## CONCLUSION

This study showed that many factors have an influence on giving antenatal care for pregnant women.

These factors play very important role in this process and we study the effect of antenatal care on the health of pregnant women.

1. The study showed that education of women play role in the quality of services. age of pregnant women also play a role in this respect.
2. Not all the women attending antenatal clinic for postnatal checking.
3. Hematinic supplementation is very good 100%. so they were all treated to prevent anaemia during pregnancy.
4. The quality of the service extended by the primary health center in a new Baqubah city was a good center. i.e 2/3 of women had antenatal visits in the first trimester of pregnancy.

## RECOMMENDATIONS

I suggest the following recommendations:-

1. Providing family planning services is an important thing to overcome the problem of unwanted pregnancy.
2. Health education for prenatal care, regular attendance of pregnant women to maternal and child health care center.
3. We must encourage adequacy of antenatal care by primary health care.
4. The record system should be done in scientific way in order to have better vital statistics.
5. There should be continuous research and evaluation of antenatal care services.

## REFERNCES

1. Amal, Diploma dissertation of community medicine, university of Baghdad. patterns of antenatal care in Sheick Omar health center and related factors, 1993.
2. Mohammed, impartial fulfilment of the requirements for the master's degree in public health, 1979.
3. Unicef, women and health 1211, Geneva, 1984; 20: 6,13, 49, 50.
4. WHO, strengthening maternal and child health program through primary health care, WHO EMRO technical publication, 1991; 18: 23–27.
5. Jawadi: MCH responsibilities and the effect on the promotion of community health. Dissertation of Diploma community of university of Baghdad, 1973.
6. Klein L. Nonregistered obstetric patient. *Am. J. Obstet Gynaco*, 1971; 110: 795–802.
7. Kaliszer and Kidd, same factors affecting attendance at antenatal clinics, *soc. Sci, med.*, 1981; 15: 421–424.
8. Hubret B, Blonedel B and Kaminskim. contribution of specialists to antenatal care infrance impact on level of care during pregnancy and delivery *J. Epid. And comm. health*, 1987; 41: 321 – 328.
9. Moore T.R, origel, w. key and Resnik R. The perinatal and economic impact of prenatal care in alow socio economic population. *Am. J. obstet, Gynaco*, 1986; 154: 29 – 33.
10. Gardiner. Motivation for obstetric care, *obst. and Gynaco*, 1969; 33: 306–312.
11. Russel I – T. British patients choice of care for Minor injury. *proc. Am. Statist. Ass.*, 1977; 548: 977.
12. Unicef, improving of maternal health in developing country. 1211 Geneve, 1984; 20: 1, 5, 6, 7, 18, 19, 22.
13. Arias, preterm labour. High risk during pregnancy and delivery. The c.v mosby company, 11830 westline industrial drive, st. Louis, mission, 1984; 63146: 37, 61.
14. Poland, M.L ager, Jw and olson J.M. Brriers to receiving adequate prenatal care *Am. J, obstel, Gynaco*, 1987; 157: 297–303.
15. Salim. is areduction of routin antenatal care Justifiable. M. sc. Dissertation, community health, university of Dublin – Trinity College, 1981.